
No.



200200132

# THE UNITED SHAMES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME;

American Takii, Inc.

MICCUS, THERE HAS BEEN PRESENTED TO THE

### Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC SEPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE HT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR RING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE PURPOSE, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT AT THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

**LETTUCE** 

'Dallas'

In Testimonn Thereof, I have hereunto set my hand and caused the seal of the Hant Haristy Itotection Office to be affixed at the City of Washington, D.C. this twenty-fifth day of August, in the year two thousand and five.

Allori

genje

Commissioner

Plant Variety Protection Office Agricultural Markoting Servico erctary of Agriculture

# Exhibit A - Origin and Breeding History

	4		æ	v	'Dallas' (Letuce D (6t: [/28]2005)
	21 22 23 24 mix	3 23 28	23 24 mix 25	21 mix	21 23 mix 24
95 S9	1		2 -	38	- 94
94 S8	1 -	3 - 22 22 23 23 23 23 23 23 23 23 23 23 23	<u>25 24</u>	31 -	36 - 12 12 36 - 3
Fixed Stage 93 S7	1- 21	<u>77 33 1</u>		20 - 1	6, 4 to 1, 2, 2, 4,
92 S6		-			e 4 æ 0 v &
91 S5		- 7			15 - 2 2 3
90 S4		244 - 1 2 3	<del>-</del>	·	284 - 1 2
89 83		1019 - 2 3 4	7607	<del>- 2</del>	1036 - 3
88 S2	- 2	818 - 4			·
87 S1	<u> </u>	509 - 5 6 7 8			
86 F1	2 - 23 23 24 24 24 24 24 24 24 24 24 24 24 24 24				
Year 85 Generation	Red Coach 74' X = 'Cisco				

**Dailos' (bt: //2/1006)**The original parents of Extrace D' are 'Red Coach 74, a Vanguard type, and 'Cisco,' a Takii original Salinas type variety.

Pallas'

Estimos Procame a variety at the S8 (F9) stage in 1994.

The fixed generation stage is S7 (F8) in 1993.

The selection criteria:

1. Head Size - to become large

- 2. Head Color to become dark green
  - 3. Head Shape to stay round

"Dallas" (bt:1/28/2005)
The color of Estimate is darker than the color of the original parent 'Red Coach 74.'
The head size of Estimate is larger than those of the original parent 'Cisco.'

The whole procedure has been processed with conventional breeding methods.

- 1. Grow trials.
- 2. Selection and grow plants in greenhouse.
  - 3. Collect seed.

'Dallas'
| Dallas | Dallas | Dallas | Dallas been observed for 6 generations of reproduction and seed increase and is stable and uniform. There are no variants except for those that would normally be expected due to the environment.

### **Exhibit B - Statement of Distinctness**

1) The most similar existing variety to 'Lettuce D' is 'Legacy.'

(bt://28/2005)

2) The characters that distinguish 'Lettuce D' from 'Legacy' are; spread of frame leaves, head weight, head diameter, core diameter, and core height.

'Dallas' (bt: 1/28/2005)
3) 'Lettuce D' is most similar to 'Legacy'; however, 'Lettuce D' has a larger spread of frame leaves than 'Legacy' (50.2 vs. 43.3 cm respectively).

'Dallas' (bt:1/28/2005)
'Lettuce D' is most similar to 'Legacy'; however, 'Lettuce D' has a greater head weight than 'Dallas' (bt:1/28/2005) 'Legacy' (777.5 vs. 512.5 g respectively).

'Dallas' (bt:1/28/2005)
'Lettuce D' is most similar to 'Legacy'; however, 'Lettuce D' has a larger head diameter than 'Legacy' (17.1 vs. 14.5 cm respectively).

'Dallas'(bt: 1/28/2005)
'Lettuce D' is most similar to 'Legacy'; however, Lettuce D' has a larger core diameter than 'Legacy' (34.6 vs. 29.6 mm respectively).

'Dallas' (bt: 1/28/2005)
-Lettuce D' has a taller core height than 'Legacy' (43.0 vs. 33.2 mm respectively).

4) Please see attached statistical data.

### U. S. Department of Agriculture Agricultural Marketing Service Science and Technology Program

### **OBJECTIVE DESCRIPTION OF VARIETY** LETTUCE Lactuca sativa

HAME OF APPLICANT (S) FOR OFFICIAL USE ONLY AMERICAN TAKII, INC.
ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code) 301 Natividad Road VARIETY NAME 'Dallas' (bt: 1/28/2005) Salinas, CA 93906 <LETTULE D> (BT:4/8/2003) Place numbers in the baxes for the characters which best describe this variety. Measured data should be the mean of an appropriate number (at least 10) of a spaced plants. Royal Horticultural Society or any recognized color standard may be used to determine plant colors. The location of the test area is: Color System Used: Salinas, ROVAL HORTICULTURAL (See list of suggested check varieties page 4.) 01\*Cutting/Leaf 05-Great Lakes Group 09=Stem 02-Butterhead 06~Vanguard Group 10-Latin 03×Bibb 07=Imperial Group 11-OTHER 04=Cos or Romaine 08=Eastern (Ithaca) Group 2. SEED: COLOR HEAT DORMANC LIGHT DORMANCY 1=White (Silver Gray) 1-Light Required 2-Black (Gray Brown) 1=Susceptible 2-Not Susceptible 2- Light Not Required 3-Brown (Amber) 3. COTYLEDON TO FOURTH LEAF STAGE: NOTE: Provide a color photograph or photocopy of the fourth test from 20 day old seedling grown under optimal conditions. SHAPE OF COTYLEDONS: 2=Intermediate 3-Spatulate SHAPE OF FOURTH LEAF: 2 3 6 LENGTH/WIDTH INDEX OF FOURTH LEAF: L/W x 10 APICAL MARGIN: 1-Entire 7-Lobed 4=Moderately Dentate 8-OTHER (specify) Z=Creanate/Gnawed 5-Coarsely Dentate BASAL MARGIN: 3-Finely Dentate 6~Incised UNDULATION: 1=Flat 2+Slight 3-Medium 4 = Marked 7-Gray Green GREEN COLOR: 1=Yellow Green 3-Medium Green 5-Blue Green GROUP 143A 2-Light Green 4=Dark Green 6=Silver Green ANTHOCYANIN: 5-OTHER (specify) 1=Absent 3-Spotted DISTRIBUTION: 2-Margin Only 4-Throughout CONCENTRATION: 1-Light 3-Intense 2-Moderate **ROLLING:** 1-Absent 2-Present CUPPING: 1=Uncupped 2-Slight 3-Markedly REFLEXING:

2=Apical Margin

3-Lateral Margins

Page 1 01

1-None

Formerly Form GR-170-1 which is obsolete

FORM LS-470-1

	<u> </u>				AAAR
		LEAVES (observe harvest-ma			
	01211	MARGIN:	nature leaves which accurately shows color	and margin characteristics.	
	2	INCISION DEPTH: Ideepest penetration of	1-Absent/Shallow (Dark Green Boston the margin)	2~Moderate (Vanguard)	3-Deep (Great Lakes 65)
	2	INDENTATION: (Minest divisions of the mars	1=Entire (Dark Green Boston) 2=Shallowly Dentate (Great Lakes 65)	3-Deeply Dentate (Great Lakes 659) 4-Cranate (Vanguard)	5-OTHER (specify)
Green		UNDULATION OF TH APICAL MARGIN:	E 1-Absent/Slight (Dark Green Boston)	2=Moderate (Vanguard)	3-Strong (Great Lakes 69
GROUP 137C	3	GREEN COLOR:	1-Very Light Green (8ibb) 2-Light Green (Minetto)	3=Medium Green (Great Lakes) 4=Dark Green (Vanguard)	5-Very Dark Green 6-OTHER
		ANTHOCYANIN (grown at	or below 10 C):		
		DISTRIBUTION:	1-Absent 2-Margin Only (Big Boston)	3=Spotted [Calif. Cream Butter] 4=Throughout (Prize Head)	5-OTHER (specify)
		CONCENTRATION:	1=Light (Iceberg)	2=Moderate (Prize Head)	3-Intense (Ruby)
	2	SIZE:	1-Small	2~Medium	3=Large
	2	GLOSSINESS:	1=Oull (Vanguard)	2=Moderate (Salinas)	3-Glossy (Great Lakes)
		BLISTERING:	1=Absent/Slight (Salinas)	2=Moderate (Vanguard)	3=Strong (Prize Head)
	2	LEAF THICKNESS:	1=Thin	2=Intermediate	3=Thick
		TRICHOMES:	1=Absent (smooth)	2-Present (spiny)	
5	<del>~~~</del> \$	SPREAD OF FRAME LEAVES cm This Variety HEAD OIAMETER (market to cm This Variety	rimmed with single cop leafs:	lspecify comparison variety	
. L	_ <del></del>	cin tins variety	1-Flattened	(specify comparison variet)	· · · · · · · · · · · · · · · · · · ·
	2	HEAD SHAPE:	2-Slightly Flattened	3=Spherical 4=Elongate	5-Non-Heading 6-OTHER
	2	HEAD SIZE CLASS:	1=Small	2=Medium	3=Large
2	_ لــــــــــــــــــــــــــــــــــــ	HEAD COUNT PER CARTON			
777	1 5	HEAD WEIGHT:  9 This Variety	5 1 2,5, Legacy	lapecily comparison variety	7
	2	MEAIL SIDMMECC.	1=Loose 2=Moderate	3=Firm 4=Very Firm	
6. BUT	T (botto	m of market-trimmed head):			
		311.00	1=Slightly Concave	2=Ftat :	3-Rounded
	A	MIDRIB:	1-Flattened (Salinas)	2-Moderately Raised	3-Prominently Raised (Grea Lakes 6591
7. CORE	lstem o	/ market-trimmed head):			60.40,000.
3	5_	om Diameter at base of head			
49	6 <sub>F</sub>	latio of head diameter/core dia	meter		
4	4 <b>~</b>	ore height from base of head to m. This Variety	33 mm Legacy	(specify comparison variety)	
8. BOLT	ING (GI	ve First Water Date	j: NOTE: First Water Date is to to germinate. This can and of	e date seed first receives adequate mois ten does equal the planting date.	
12	-	umber of days from First Wate his Variety	or Date to seed stalk emergence (summer co		
	<u>3</u> B				5-Very Rapid
14	<i>7</i> 1	eight of mature seed stalk: n. This Variety	139 cm Legacy	(specify comparison variety)	

FORM LS-470-1

BOLTER LEAVES:    1-Straight   1-Enrine   2-Deniase	42	Spread of Bolter Plant for win	dest point):	Legacy	(specify comparison v	ariety)
AARGIN: 1-Entire 2-Dentals    COLOR:		BOLTER LEAVES:	1-Straight	2=Curv	red	
BOLTER HABIT:  2 ITEMMINAL INFLORESCENCE: 1-Absent 2-Present  2 LATERAL SHOOTS: 1-Absent 2-Present  2 BASAL SIDE SHOOTS: 1-Absent 2-Present  3 MATURITY territores of harvest-mature head formation]:  NOTE: Complete this section for at less one sensor.  SEASON Applic U Foldays Cheek U Foldays  Spring Summer 6 5 6 7 Legacy  Winter  Give planting date(s), and location(s):  Spring Summer Seeded 7/23/01 in Salinas, CA  Fall Seeded 8/4/01 in Salinas, CA  Winter  U First water date to harvest. U Fill in check variety name on the appropriate line.  10. ADAPTATION:  PRIMARY REGIONS OF ADAPTION lessed and proven adapted): (0-Not tested 1+Not Adapted)  2 Southwest (Calif., Arz. deset)  2 Southwest (Calif., Arz. deset)  2 Southwest (Calif., Arz. deset)  2 Southwest (Calif., Spring Seas Southwest + West Coast  2 Summer larce Southwest + West Coast  2 Summer larce Southwest + West Coast  3 Southwest (real of the season)  4 Winter (seas Southwest)  4 Winter (seas Southwest)  5 Summer larce Southwest + West Coast  C Orner (seas Southwest)  C REENHOUSE: 0-Not tested 1-Not Adapted 2-Adapted		MARGIN:	1=Entire	2-Den	1316	
TERMINAL INFORMESCENCE: 1-Absent 2-Present    2	2	COLOR:	→ Light Green	2-Med	ium Green	3+Dark Green
INFLORESCENCE:    2		BOLTER HABIT:	,			
ASAL SIDE SHOOTS: 1-Absent 2-Present	2		T-Absent	2-Press	ent	<i>'</i>
9. MATURITY (retilines of harvest-mediar head formation):  NOTE: Complete this section for at less one sevice.  SEASON Applic U od days Check U od days  Spring  Summer 65 67 Legacy  Winter  Give planting date(s), and location(s):  Spring  Summer Seeded 7/23/01 in Salimas CA  Fall Seeded 8/4/01 in Salimas CA  Winter  U First water date to harvest. Uffit in check variety name on the appropriate lina.  10. ADAPTATION:  PRIMARY REGIONS OF ADAPTION (tested and proven adapted): (0-Not tested 1=Not Adapted 2=Adapted)  PRIMARY REGIONS OF ADAPTION (tested and proven adapted): (0-Not tested 1=Not Adapted 2=Adapted)  SSALON:  S	2		1-Absent	2-Prese	ent	
SEASON Applic II For days Check II Foldays CHECK VARIETY II  Spring Summer 665 67 Legacy  Fall 779 81 Legacy  Winter Seeded 7/23/01 in Salinas CA  Fall Seeded 8/4/01 in Salinas CA  Winter II First water date to hervest. If it is check variety asms on the appropriate line.  10. ADAPTATION:  PRIMARY REGIONS OF ADAPTION (tested and proven adapted): (0-Not tested 1=Not Adapted 2-Adapted)  2 Southwest (Collif., Aris. desert)	2	BASAL SIDE SHOOTS:	1-Absent	2=Presi	ent	
SEASON Applic I Fold days Check I Fold days Check VARIETY I Spring  Summer 655 67 Legacy  Winter  Give planting date(s), and location(s):  Spring  Summer Seeded 7/23/01 in Salinas, CA  Fall Seeded 8/4/01 in Salinas, CA  Winter  I) First water date to harvest. If it in chack variety name on the appropriate line.  10. ADAPTATION:  PRIMARY REGIONS OF ADAPTION (tested and proven adapted): (0-Not tested 1-Not Adapted 2-Adapted)  2 Southwest (Colif., Ariz. desert)	9. MATURITY	learliness of harvest-mature h	ead formation):			
SEASON Applic Design Control of the Secret Control of Spring Summer 665 67 Cegacy  Winter 79 81 Legacy  Winter Seeded 7/23/01 in Salinas, CA  Fall Seeded 8/4/01 in Salinas, CA  Winter Definite water date to harvest. 3/Fill in check variety name on the appropriate line.  10. ADAPTATION:  PRIMARY REGIONS OF ADAPTION Itested and proven adapted): (0-Not tested 1-Not Adapted 2-Adapted)  2 Southwest (Calif., Aris. desert) 2 West Coast 0 Northeast  O Northeastal 2 OTHER Japan  SEASON:  2 Southwest (Calif. Southwest + West Coast 2 Fall (ares Southwest + West Coast 2 Summer (area Southwest + West Coast 1 Winter (area 1-Not Adapted 2-Adapted 2-Adap	NOTE: 0	·			CHECK VARIETY	2/
Summer  Fall  79  81  Legacy  Winter  Give planting date(s), and location(s):  Spring  Summer Seeded 7/23/01 in Salinas, CA  Fall Seeded 8/4/01 in Salinas, CA  Winter  UFirst water date to harvest. 2/Fill in check variety name on the appropriate lina.  10. ADAPTATION:  PRIMARY REGIONS OF ADAPTION (tested and proven adapted): (0-Not tested 1-Not Adapted 2-Adapted)  2 Southwest (Calif., Arts. desert)  2 West Coast  O Northcentral  SEASON:  2 Spring (area Southwest + West Coast  2 Summer (area Southwest + West Coast  1 Winter (area  O GREENHOUSE: 0-Not tested 1-Not Adapted 2-Adapted	SEAS	SON Applic 1 *of day	Check 1/ #of days		CHECK VARIETY	
Give planting date(s), and location(s):  Spring  Summer Seeded 7/23/01 in Salinas, CA  Fall Seeded 8/4/01 in Salinas, CA  Winter  1/ First water date to harvest.  2/ Fill in check variety name on the appropriate line.  10. ADAPTATION:  PRIMARY REGIONS OF ADAPTION (tested and proven adapted): (0-Not tested 1-Not Adapted 2-Adapted)  2 Southwest (Calif., Aris. desert)  O Northcentral  O Southwest  SEASON:  ONTHER  SEASON:  SEASON:  SEASON:  ONTHER  SEASON:  SEASON:  ONTHER  SEASON:  SEASON:  ONTHER  SEASON:	Spring	,				· · · · · · · · · · · · · · · · · · ·
Give planting date(s), and location(s):  Spring  Summer Seeded 7/23/01 in Salinas, CA  Fall Seeded 8/4/01 in Salinas, CA  Winter  1/ First water date to harvest.  2/ Fill in check variety name on the appropriate line.  10. ADAPTATION:  PRIMARY REGIONS OF ADAPTION (tested and proven adapted): (0-Not tested 1-Not Adapted 2-Adapted)  2 Southwest (Calif., Aris. desert)  O Northcentral  O Southwest  SEASON:  ONTHER  SEASON:  SEASON:  SEASON:  ONTHER  SEASON:  SEASON:  ONTHER  SEASON:  SEASON:  ONTHER  SEASON:		[[.5]	67	Legary	•	
Give planting date(s), and location(s):  Spring  Summer Seeded 7/23/01 in Salinas, CA  Fall Seeded 8/4/01 in Salinas, CA  Winter  If First water date to harvest. 2/Fill in check variety name on the appropriate line.  10. ADAPTATION:  PRIMARY REGIONS OF ADAPTION (tested and proven adapted): (0-Not tested 1-Not Adapted 2-Adapted)  2 Southwest (Colif., Ariz. desert)  O Northcentral  SEASON:  SEASON:	Summ					
Give planting date(s), and location(s):  Spring  Summer Seeded 7/23/01 in Salinas CA  Fall Seeded 8/4/01 in Salinas CA  Winter  Uffirst water date to harvest. 2/Fill in check variety name on the appropriate line.  10. ADAPTATION:  PRIMARY REGIONS OF ADAPTION (tested and proven adapted): (0-Not tested 1-Not Adapted 2-Adapted)  2 Southwest (Collif., Ariz., desert) 2 West Coast 0 Northeast  O Northcentral 0 Southwest 2 OTHER JAPAN  SEASON: 2 Spring (area Southwest + West Coast 1 Winter (area	Fall	7 9	81	Legacy		
Summer Seeded 7/23/01 in Salinas, CA  Fall Seeded 8/4/01 in Salinas, CA  Winter  1/ First water date to harvest. 2/ Fill in check variety name on the appropriate time.  10. ADAPTATION:  PRIMARY REGIONS OF ADAPTION (tested and proven adapted): (0-Not tested 1-Not Adapted 2-Adapted)  2 Southwest (Calif., Ariz. desert)  Northcentral  SEASON:  SEASON:  2 Spring (area Southwest + West Coast 2 Fall (area Southwest + West Coast 3 Fall (area Southwest + West Coast 4 Fall (area Southwest + West Coast 4 Fall (area Southwest + West Coast 5 Fall (area Southwest + West Coast 6 Fall (area Southwest + West Coast 7 Fall (area Southwest 7 Fall (area Southwest 7 Fall (area Southwest 7 Fa	Winte					
Summer Seeded 7/23/01 in Salinas, CA  Fall Seeded 8/4/01 in Salinas, CA  Winter  1/ First water date to harvest. 2/ Fill in check variety name on the appropriate time.  10. ADAPTATION:  PRIMARY REGIONS OF ADAPTION (tested and proven adapted): (0-Not tested 1-Not Adapted 2-Adapted)  2 Southwest (Calif., Ariz. desert)  Northcentral  SEASON:  SEASON:  2 Spring (area Southwest + West Coast 2 Fall (area Southwest + West Coast 3 Fall (area Southwest + West Coast 4 Fall (area Southwest + West Coast 4 Fall (area Southwest + West Coast 5 Fall (area Southwest + West Coast 6 Fall (area Southwest + West Coast 7 Fall (area Southwest 7 Fall (area Southwest 7 Fall (area Southwest 7 Fa	Give planting	date(s), and location(s):		· · · · · · · · · · · · · · · · · · ·		
Summer Seeded 7/23/01 in Salinas CA  Fall Seeded 8/4/01 in Salinas, CA  Winter  1/ First water date to harvest. 2/Fill in check variety name on the appropriate line.  10. ADAPTATION:  PRIMARY REGIONS OF ADAPTION (tessed and proven adapted): (0-Not tested 1=Not Adapted 2=Adapted)  2 Southwest (Calif., Ariz. desert) 2 West Coast 0 Northeast  0 Northcentral 0 Southwest 2 OTHER Japan  SEASON: 2 Spring (area Southwest + West Coast 2 Fall (area Southwest + West Coast 3 Winter (area 1 Winter (area 1 - Not Adapted 2-Adapted)  0 GREENHOUSE: 0-Not tested 1=Not Adapted 2-Adapted			~			
Winter  If First water date to harvest.  2/Fill in check variety name on the appropriate line.  10. ADAPTATION:  PRIMARY REGIONS OF ADAPTION (tested and proven adapted): (0-Not tested 1=Not Adapted 2=Adapted)  2 Southwest (Calif., Ariz. desert)  Northcentral  SEASON:  SEAS			2/	Salinas CA		
Winter  If First water date to harvest.  If Fill in check variety name on the appropriate line.  10. ADAPTATION:  PRIMARY REGIONS OF ADAPTION lessed and proven adapted:  Onortheast  Onortheast  Onortheast  SEASON:  Seas	Summ				<u> </u>	
10. ADAPTATION:  PRIMARY REGIONS OF ADAPTION (tested and proven adapted): (0-Not tested 1-Not Adapted 2-Adapted)  2 Southwest (Calif., Ariz. desert)  O Northcentral  SEASON:  SEASON:  Septing (area Southwest + West Coast  C Southwest + West Coast  D Southwest + West Coast  C Southwest + West Coast  D Northcentral  SEASON:  SEASON:  O Northcentral  O Southwest + West Coast  O Northcentral  O Northcentral  SEASON:  O Northcentral  O Northcentral  SEASON:  O Northcentral  O Northcen	Fall	Seeded 8/4	f/01 in S	SAliNAS, CA		
10. ADAPTATION:  PRIMARY REGIONS OF ADAPTION (tested and proven adapted): (0-Not tested 1-Not Adapted 2-Adapted)  2 Southwest (Calif., Ariz. descrit)  O Northcentral  SEASON:  2 Spring (area Southwest + West Coast  2 Spring (area Southwest + West Coast  Winter (area Southwest + West Coast  O GREENHOUSE: 0-Not tested 1-Not Adapted 2-Adapted			2/Fill in check yar	lety name on the appropris	ge line.	
PRIMARY REGIONS OF ADAPTION (tested and proven adapted): (0=Not tested 1=Not Adapted 2=Adapted)  2 Southwest (Calif., Ariz. desert)  2 West Coast  D Northcentral  SEASON:  2 Spring (area Southwest + West Coast  D Southwest + West Coast  D Southwest + West Coast  D Northcentral  SEASON:  2 Spring (area Southwest + West Coast  D Northcentral  SEASON:  1 Fall (area Southwest + West Coast  D Winter (area Southwest + West Coast  D Northcentral  SEASON:  1 Fall (area Southwest + West Coast  D Northcentral  2 Fall (area Southwest + West Coast  D Northcentral  2 Fall (area Southwest + West Coast  D Northcentral  2 Fall (area Southwest + West Coast  D Northcentral  3 Fall (area Southwest + West Coast  D Northcentral  2 Fall (area Southwest + West Coast  D Northcentral  2 Fall (area Southwest + West Coast  D Northcentral  2 Fall (area Southwest + West Coast  D Northcentral  2 Fall (area Southwest + West Coast  D Northcentral  2 Fall (area Southwest + West Coast  D Northcentral  2 Fall (area Southwest + West Coast  D Northcentral	<del></del>		3			
O Northcentral  SEASON:  SEASO			APTION (tested and pro	ven adapted): (0=Not	tested 1=Not Ada	opted 2=Adapted
2 Summer farea Southwest + West Coast    Winter farea   Yest Coast   Winter farea   Yest Coast	2	Southwest (Calif., Ariz, desert)	West Co.	ast Northea	ist	
2 Summer farea Southwest + West Coast    Winter farea   Yest Coast   Winter farea   Yest Coast	0	Northcentral	O Southea	st 2 OTHER	JAPAN	
Z Summer larea Southwest + West Coast  Winter larea  1-Not Adapted  2-Adapted		SEASON:  2 Spring (area Southwe:	it + West Coast	<b>2</b> Fall (are	southwest + Wes	st Coast,
GREENHOUSE: 0-Not tested 1-Not Adapted 2-Adapted	į		est + West GAS	+ Winter	(area	1
7-90ID			0-Nat tested	1-Not /	Adapted	2-Adapted
3 SOIL TYPE: 1-Mineral 2-Organic 3-00		SOIL TYPE:	1-Mineral	2-01981	nic	3-8oth

200200132

·	
11. DISEASES AND STRESS REACTIONS (0=Not tested; 1=Suscept	tible; 2*Intermediate; 3=Resistant; 4*Highly resistant; 5=Tolerant);
VIRUS	FUNGAL/BACTERIAL
<b>∏</b> Big Vein	Corky Root Rot (Pythium Root Rot)
Lettuce Mosaic	Downy Mildew (Races)
Oucumber Mosaic	2 Powdery Mildew
Sroad Bean Wilt	2 Sclerotinia Rot
Turnip Mosaic	2 Bacterial Soft Rot (Pseudomonas spp. & others)
Seet Western Yellows	O Botrytis (Gray Mold)
Collection Vellows	OTHER
Other Virus	<u>.</u>
INSECTS	PHYSIOLOGICAL/STRESS
INSECTS  Cabbage Loopers	Z Tipburn O Salt
Root Aphids	Heat 2 Brown Rib (Rib Discoloration, Rib Blig
Green Peach Aphid	2 Drought OTHER
Other Insect	Z Cold
2005	. UA DVSST
2 Pink Rib	HARVEST  Internal Rib Necrosis (Blackheart, Gray Rib, Gray Streak)
Russet Sporting	Brown Stain
2 Rusty Brown Discoloration	
2. BIOCHEMICAL OR ELECTROPHORETIC MARKERS:	•
None	
13. COMMENTS:	
Sisten line of 'Legacy' which is A	popular commercial Salinas variety.
the state of the s	D' is Lapari improved.
Sister line of 'Legacy' which is A 'Legacy' does not have PVP. 'Lettuce	D is Legacy improved.

SUGGESTED CHECK VARIETIES

### TYPE

- 1) CUTTING/LEAF
  2) BUTTERHEAD
  3) BIBB
  4) COS, OR ROMAINE
  5) GREAT LAKES GROUP
  6) VANGUARD GROUP
  7) IMPERIAL GROUP
  8) EASTERN GROUP
  9) STEM
  10) LATIN

### CHECK VARIETY

SALAD BOWL DARK GREEN BOSTON BIBB PARRIS ISLAND GREAT LAKES 659-700 VANGUARD VIVA ITHAÇA CELTUCE MATCHLESS

200200132

### Trial 1

Location: Frank McFadden Ranch Lot 3F, Salinas, CA

Planting Date: 7/23/01 Evaluation Date: 9/26/01

Plot Size: Four 20 ft. plots with 172 plants total

Head Weight (g)

Dead Xveight (d)					
<u> </u>	Legacy				
1025	550				
875	425				
850	625				
575	450				
725	575				
850	600				
925	475				
750	400				
650	600				
950	400				

Data Summary

	Data Sulfillary					
Ŀ	:428/105)	Lettuce D	Legacy	Total		
	Ν	10	10	20		
	-Σ X	8175	5100	13275		
Ì	-Mean	817.5	510	663.75		
	-Σ X <sup>2</sup>	6860625	2672500	9533125		
	Variance	19729.17	7944.444	37991.78		
ı	Std. Dev.	140.4606	89.1316	194.9148		
	Std. Err.	44.4175	28.1859	43.5843		

**ANOVA Summary** 

7.110 171 Guillinary						
Source	SS	df	MS	F	Р	
Treatment	472781.3	1	472781.3	34.17	<.0001	
Error	249062.5	18	13836.81			
Total	721843.8	19				

F Distribution

Table Value of F <sub>.05</sub>	4.41
Table Value of F <sub>.01</sub>	8.29
Computed F Ratio	34.17

Computed F ratio > table value of F<sub>.01</sub> so the means are highly significantly different at 1%.

### Trial 2

Location: Secondo Farms Lot 11, Salinas, CA

Planting Date: 8/4/01 Evaluation Date: 10/19/01

Plot Size: Four 100 ft. plots with 864 plants total

Head, Weight (g)

<del>L'offlice D</del>	Legacy
700	600
850	450
775	550
750	450
675	525
575	575
750	450
675	475
800	500
825	575
•	

Data Summary

1/28/2005)	Lettuce D	Legacy	Total
N	10	10	20
-Σ X	7375	5150	12525
-Mean	737.5	515	626.25
-Σ X <sup>2</sup>	5500625	2682500	8183125
Variance	6840.278	3361.111	17860.2
Std. Dev.	82.706	57.9751	133.642
Std. Err.	26.1539	18.3333	29.8833

**ANOVA Summary** 

Source	SS	df	MS	F	Р
Treatment	247531.3	1	247531.3	48.53	<.0001
Error	91812.5	18	5100.69	****	
Total	339343.8	19			

F Distribution

Table Value of F <sub>.05</sub>	4.41
Table Value of F <sub>.01</sub>	8.29
Computed F Ratio	48.53

Computed F ratio > table value of F<sub>.01</sub> so the means are highly significantly different at 1%.

Average Mean of Trial 1 and Trial 2

Leftuce D Legacy 777.5 512.5

### Trial 1

Location: Frank McFadden Ranch Lot 3F, Salinas, CA

Planting Date: 7/23/01 Evaluation Date: 9/26/01

Plot Size: Four 20 ft. plots with 172 plants total

Spread of Frame Leaves (cm)

	Spread of	Frame Leav
J	<del>Collins D</del>	Legacy
	53.0	48.0
Ī	50.0	47.0
	51.0	45.5
	52.0	37.5
	51.0	45.5
	51.0	47.0
I	47.5	42.5
l	47.0	40.5
ı	47.5	44.5
I	51.0	46.5
•		

**Data Summary** 

Bata Cammary					
t 1/28/05) .	Lettuce D	Legacy	Total		
N	10	10	20		
-Σ X	501	444.5	945.5		
-Mean	50.1	44.45	47.275		
-Σ X <sup>2</sup>	25138.5	19857.75	44996.25		
Variance	4.2667	11.0806	15.6704		
Std. Dev.	2.0656	3.3287	3.9586		
Std. Err.	0.6532	1.0526	0.8852		
	N -Σ X -Mean -Σ X <sup>2</sup> Variance Std. Dev.	N 10 -Σ X 501 -Mean 50.1 -Σ X² 25138.5 Variance 4.2667 Std. Dev. 2.0656	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		

**ANOVA Summary** 

Source	SS	df	MS	F	Р	
Treatment	159.61	1	159.61	20.81	0.000242	
Error	138.13	18	7.67			
Total	297.74	19				

F Distribution

Table Value of F <sub>.05</sub>	4.41
Table Value of F <sub>.01</sub>	8.29
Computed F Ratio	20.81

Computed F ratio > table value of F<sub>.01</sub> so the means are highly significantly different at 1%.

### Trial 2

Location: Secondo Farms Lot 11, Salinas, CA

Planting Date: 8/4/01 Evaluation Date: 10/19/01

Plot Size: Four 100 ft. plots with 864 plants total

Spread of Frame Leaves (cm)

į	<del>Lettuco D</del>	Legacy
	52.0	46.0
	51.0	40.5
	51.5	40.0
	48.0	42.0
	47.5	39.5
	49.0	42.5
	52.0	41.0
	52.5	42.0
İ	46.0	43.0
Į	53.0	45.5

Data Summary 105 (1511/28/2005) | Lettuce E

Data Carrier 106						
2005)	-Eettuce-D-	Legacy	Total			
N	10	10	20			
X	502.5	422	924.5			
ean	50.25	42.2	46.225			
$X^2$	25303.75	17851	43154.75			
ance	5.9028	4.7333	22.0914			
Dev.	2.4296	2.1756	4.7002			
Err.	0.7683	0.688	1.051			
	ean X <sup>2</sup> ance Dev.	Property         Property           N         10           CX         502.5           ean         50.25           X²         25303.75           ance         5.9028           Dev.         2.4296	A005         Eetfuce D         Legacy           N         10         10           CX         502.5         422           ean         50.25         42.2           X²         25303.75         17851           ance         5.9028         4.7333           Dev.         2.4296         2.1756			

**ANOVA Summary** 

,					
Source	SS	df	MS	F	Р
Treatment		1	324.01	60.9	<.0001
Error	95.73		5.32		
Total	419.74	19			

F Distribution

Table Value of F <sub>.05</sub>	4.41
Table Value of F <sub>.01</sub>	8.29
Computed F Ratio	60.9

Computed F ratio > table value of  $F_{.01}$  so the means are highly significantly different at 1%.

) <del>Lettice D</del>	Legacy
50.175	43.325

### Trial 1

Location: Frank McFadden Ranch Lot 3F, Salinas, CA

Planting Date: 7/23/01 Evaluation Date: 9/26/01

Plot Size: Four 20 ft. plots with 172 plants total

	Head Dian	neter (cm)
j	Lettuce D	Legacy
1	18.2	16.0
	17.3	15.7
İ	17.2	14.4
	17.3	15.0
	16.0	14.5
	16.5	15.5
	18.5	15.0
l	18.2	13.5
	17.0	14.0
ĺ	15.0	15.2

Data Summary

Data Summary					
:1/28/2005	Lettuce D	Legacy	Total		
N	10	10	20		
-Σ X	171.2	148.8	320		
-Mean	17.12	14.88	16		
-Σ X <sup>2</sup>	2941.4	2219.64	5161.04		
Variance	1.1618	0.6107	2.16		
Std. Dev.	1.0779	0.7815	1.4697		
Std. Err.	0.3408	0.2471	0.3286		
	N -Σ X -Mean -Σ X² Variance Std. Dev.	N 10 -Σ X 171.2 -Mean 17.12 -Σ X² 2941.4 Variance 1.1618 Std. Dev. 1.0779	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$		

**ANOVA Summary** 

Source	SS	df	MS	F	Р
Treatment	25.09	1	25.09	28.19	<.0001
Error	15.95	18	0.89		
Total	41.04	19			

F Distribution

Table Value of F <sub>.05</sub>	4.41
Table Value of F <sub>.01</sub>	8.29
Computed F Ratio	28.19

Computed F ratio > table value of F<sub>.01</sub> so the means are highly significantly different at 1%.

Location: Secondo Farms Lot 11, Salinas, CA

Planting Date: 8/4/01 Evaluation Date: 10/19/01

Plot Size: Four 100 ft. plots with 864 plants total

13.5

14.0

14.0

Head Diameter (cm)

Legacy 15.1 13.9 17.5 15.0 15.5 15.0 13.0 19.7 14.5 14.0 16.5 14.5 17.0 13.5

**Data Summary** 

I	t:1/28/105)	Lettuce D	Legacy	Total
	N	Dallas 10	10	20
	-Σ X	170.3	140.4	310.7
	-Mean	17.03	14.04	15.535
	$-\Sigma X^2$	2927.35	1974.96	4902.31
	Variance	3.0157	0.416	3.9782
	Std. Dev.	1.7366	0.645	1.9945
	Std. Err.	0.5492	0.204	0.446

**ANOVA Summary** 

19.5

18.0

17.0

Source	SS	df	MS	F	Р
Treatment	44.7	1	44.7	25.99	<.0001
Error	30.89	18	1.72		
Total	75.59	19			

**F** Distribution

Table Value of F <sub>.05</sub>	4.41
Table Value of F <sub>.01</sub>	8.29
Computed F Ratio	25.99

Computed F ratio > table value of F<sub>.01</sub> so the means are highly significantly different at 1%.

200200132

### Trial 1

Location: Frank McFadden Ranch Lot 3F, Salinas, CA

Planting Date: 7/23/01 Evaluation Date: 9/26/01

Plot Size: Four 20 ft. plots with 172 plants total

### Core Diameter (mm)

	COLE DIGIL	Core Diameter (min)			
	<del>Lettuce D</del>	Legacy			
(bt:1/28/	2 <b>005</b> ) 37	30			
·	35	27			
	35	33			
	33	30			
	33	31			
	35	31			
	37	30			
	33	26			
	35	29			
	36	27			

### Data Summary

	Data Sullillary				
(bt: 1/28/2005)		<del>Lettuce D</del>	Legacy	Total	
	N	10 Danies	10	20	
	-Σ X	349	294	643	
	-Mean	34.9	29.4	32.15	
	-Σ X <sup>2</sup>	12201	8686	20887	
	Variance	2.3222	4.7111	11.2921	
	Std. Dev.	1.5239	2.1705	3.3604	
	Std. Err.	0.4819	0.6864	0.7514	

### **ANOVA Summary**

7,010 171 00					
Source	SS	df	MS	F	Р
Treatment	151.25	1	151.25	42.97	<.0001
Error	63.3	18	3.52		
Total	214.55	19			

### F Distribution

Table Value of F <sub>.05</sub>	4.41
Table Value of F <sub>.01</sub>	8.29
Computed F Ratio	42.97

Computed F ratio > table value of  $F_{.01}$  so the means are highly significantly different at 1%.

### Trial 2

Location: Secondo Farms Lot 11, Salinas, CA

Planting Date: 8/4/01 Evaluation Date: 10/19/01

Legacy

31

28

31

Plot Size: Four 100 ft. plots with 864 plants total

### Core Diameter (mm)

(bt:1/28/2	34	- 3
• •	36	2
	35	3
	36	2
	36	2
	29	31
	33	2

**Data Summary** 

	Data Gaimilary				
(bt:1/2 <i>8/2005</i> ) N		<del>Lettuce D</del>	Legacy	Total	
		Pallas 10	10	20	
	-Σ X	342	298	640	
	-Mean	34.2	29.8	32	
	-Σ X <sup>2</sup>	11738	8912	20650	
	Variance	4.6222	3.5111	8.9474	
	Std. Dev.	2.1499	1.8738	2.9912	
	Std. Err.	0.6799	0.5925	0.6689	

### **ANOVA Summary**

33

35

35

	Source	SS	df	MS	F	Р
	Treatment	96.8	1	96.8	23.78	0.000121
	Error	73.2	18	4.07		
i	Total	170	19			

### F Distribution

Table Value of F <sub>.05</sub>	4.41
Table Value of F <sub>.01</sub>	8.29
Computed F Ratio	23.78

Computed F ratio > table value of  $F_{.01}$  so the means are highly significantly different at 1%.

(bt:1/204	Lettuce D	Legacy
100/70	<b>95)</b> 34.55	29.6

### Trial 1

Location: Frank McFadden Ranch Lot 3F, Salinas, CA

Planting Date: 7/23/01 Evaluation Date: 9/26/01

Plot Size: Four 20 ft. plots with 172 plants total

Core Height (mm)

		(0,,,,,,
,	Lettuce D	Legacy
(bt:1/28/	2005) 47	33
,	44	25
	42	35
	42	40
	45	36
	35	33
	53	28
	43	31
	35	37
	44	33

Data Summary

	Data Summary				
(bt	:1/28/2005).	<del>Lettuce D</del>	Legacy	Total	
	Ν	Dail 45 10	10	20	
	-Σ X	430	331	761	
	-Mean	43	33.1	38.05	
	-Σ X <sup>2</sup>	18742	11127	29869	
	Variance	28	18.9889	48.05	
	Std. Dev.	5.2915	4.3576	6.9318	
	Std. Err.	1.6733	1.378	1.55	

**ANOVA Summary** 

	,				
Source	SS	df	MS	F	Р
Treatment	490.05	1	490.05	20.86	0.000239
Error	422.9	18	23.49		
Total	912.95	19			

F Distribution

Table Value of F <sub>.05</sub>	4.41
Table Value of F <sub>.01</sub>	8.29
Computed F Ratio	20.86

Computed F ratio > table value of F<sub>.01</sub> so the means are highly significantly different at 1%.

### Trial 2

Location: Secondo Farms Lot 11, Salinas, CA

Planting Date: 8/4/01 Evaluation Date: 10/19/01

Plot Size: Four 100 ft. plots with 864 plants total

Core Height (mm)

(bt:1/28/	2005) 47	4
	37	2
	50	3
	44	3
	40	2
	32	3

40	29
32	38
48	25
41	34
42	31
. 49	34

Data Summary

_	Pata Suilli	144 DE 10 C		
Gt.	/28/2005)	Lettuce D	Legacy	Total
	N	10	10	20
	-Σ X	430	333	763
	-Mean	43	33.3	38.15
	-Σ X <sup>2</sup>	18788	11445	30233
	Variance	33.1111	39.5667	59.1868
	Std. Dev.	5.7542	6.2902	7.6933
	Std. Err.	1.8196	1.9891	1.7203

**ANOVA Summary** 

Source	SS	df	MS	F	Р
Treatment	470.45	1	470.45	12.95	0.002054
Error	654.1	18	36.34		
Total	1124.55	19			

F Distribution

Table Value of F <sub>.05</sub>	4.41
Table Value of F <sub>.01</sub>	8.29
Computed F Ratio	12.95

Computed F ratio > table value of F<sub>.01</sub> so the means are highly significantly different at 1%.

	<b>CHOOSE</b>	77.1	cuii oi	TTIG
bt:1/20	Lettu	a. G	Lega	су
DU: YOU	10005)	43	3	3.2

### Trial 1

Location: Frank McFadden Ranch Lot 3F, Salinas, CA

Planting Date: 7/23/01 Evaluation Date: 9/26/01

Plot Size: Four 20 ft. plots with 172 plants total

### Height of Mature Seed Stalk (cm)

	neight of Mature Seet		
	- <del>Lettuce, D</del>	Legacy	
Cbt:1/28	Dallas 35	131	
·	142	133	
	139	134	
	157	149	
	156	121	
	140	136	
	143	114	
	104	135	
	141	115	
	150	146	

Data Summary (128/2005)							
	Lettuce D Legacy						
N	10	10	20				
-Σ X	1407	1314	2721				
-Mean	140.7	131.4	136.05				
-Σ X <sup>2</sup>	199941	173906	373847				
Variance	219.5667	138.4889	192.3658				
Std. Dev.	14.8178	11.7681	13.8696				
Std. Err.	4.6858	3.7214	3.1013				

**ANOVA Summary** 

Source	SS	df	MS	F	Р		
Treatment	432.45	1	432.45	2.42	0.137202		
Error	3222.5	18	179.03				
Total	3654,95	19					

### F Distribution

Table Value of F <sub>.05</sub>	4.41
Table Value of F <sub>.01</sub>	8.29
Computed F Ratio	2.42

Computed F ratio < table values of F so the means are not significantly different.

### Trial 2

Location: Secondo Farms Lot 11, Salinas, CA

Planting Date: 8/4/01 Evaluation Date: 10/19/01

Plot Size: Four 100 ft. plots with 864 plants total

### Height of Mature Seed Stalk (cm)

Lettuce D	Legacy
34	135
145	145
142	151
146	149
127	155
147	150
145	153
151	142
146	150
147	137

Data Summary

(bt:1/28/05)	<del>Lettuce D</del>	Legacy	Total			
N	10	10	20			
-Σ X	1430	1467	2897			
-Mean	143	146.7	144.85			
-Σ X <sup>2</sup>	204950	215619	420569			
Variance	51.1111	45.5667	49.3974			
Std. Dev.	7.1492	6.7503	7.0283			
Std. Err.	2.2608	2.1346	1.5716			

**ANOVA Summary** 

Source	SS	df	MS	F	Р
Treatment	68.45	1	68.45	1.42	0.248879
Error	870.1	18	48.34		
Total	938.55	19			

F Distribution

Table Value of F <sub>.05</sub>	4.41
Table Value of F <sub>.01</sub>	8.29
Computed F Ratio	1.42

Computed F ratio < table values of F so the means are not significantly different.

-Eellise D	Legacy
141.85	139.05

### Trial 1

Location: Frank McFadden Ranch Lot 3F, Salinas, CA

Planting Date: 7/23/01 Evaluation Date: 9/26/01

Plot Size: Four 20 ft. plots with 172 plants total

### Spread of Bolter Plant (cm)

	Spread of	boiler Plan
	<del>Lettuce D</del>	Legacy
1	20 <b>5</b> ) 36	34
	32	32
	44	36
	39	36
	40	30
	43	36
	32	32
İ	30	43
	31	28
I	35	38
•		

Da	ıta	Su	ımı	ma	ry
•	- 1		_		

	Bata Cammary				
(kt. 1/28/2005)		<del>Lettuec D</del>	Legacy	Total	
-	N	10 10 m	10	20	
	-Σ X	362	345	707	
i	-Mean	36.2	34.5	35,35	
	-Σ X <sup>2</sup>	13336	12069	25405	
	Variance	25.7333	18.5	21.7132	
	Std. Dev.	5.0728	4.3012	4.6597	
	Std. Err.	1.6042	1.3601	1.0419	

### **ANOVA Summary**

,							
Source	SS	df	MS	F	Р		
Treatment	14.45	1	14.45	0.65	0.430636		
Error	398.1	18	22.12				
Total	412.55	19					

### F Distribution

Table Value of F <sub>.05</sub>	4.41
Table Value of F <sub>.01</sub>	8.29
Computed F Ratio	0.65

Computed F ratio < table values of F so the means are not significantly different.

### Trial 2

Location: Secondo Farms Lot 11, Salinas, CA

Planting Date: 8/4/01 Evaluation Date: 10/19/01

Plot Size: Four 100 ft. plots with 864 plants total

### Spread of Bolter Plant (cm)

(bt	·/a	8/2	

Legacy
39
42
41
39
42
38
39
44
44 43
41

**Data Summary** 

Ы	:: 1/28/105)	<del>Lettuce D-</del>	Legacy	Total	
	N	Dallas 10	10	20	
	-Σ X	473	408	881	
	-Mean	47.3	40.8	44.05	
ĺ	$-\Sigma X^2$	23231	16682	39913	
	Variance	95.3444	3.9556	58.1553	
	Std. Dev.	9.7644	1.9889	7.626	
	Std. Err.	3.0878	0.6289	1.7052	

### **ANOVA Summary**

Source	SS	df	MS	F	Р
Treatment	211.25	1	211.25		0.053995
Error	893.7	18	49.65		
Total	1104.95	19			

### F Distribution


Distribution	
Table Value of F <sub>.05</sub>	4.41
Table Value of F <sub>.01</sub>	8.29
Computed F Ratio	4.25

Computed F ratio < table values of F so the means are not significantly different.


Lettuce D	Legacy	
41.75	37.65	

# 200.00132


# 200200132




Fourth leaf from a 20 day old seedling of 'Lettuce D.'
'Dallas' (1/28/2005) bt)



Harvest mature leaves of 'Lettuce D.'
'Dallas' (1/28/2005) ot)



Sample of Lettuce D' from field trial. Dollars (bt: 1/38/2005)



Sample of 'Legacy' - the most similar variety to 'Lettuce D.'
Delias' (bt=1/28/2008)



Fourth leaf from 20 day old seedling of Lettuce "D" Dallos (1614 1/28/8005)

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE	The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Paperwork Reduction Act (PRA) of 1995.		
EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP		etermine if a plant variety protection 421). Information is held confidential	
1. NAME OF APPLICANT(S)  American Takii, Inc.	TEMPORARY DESIGNATION     OR EXPERIMENTAL NUMBER	3. VARIETY NAME	
MINITERIORIA IMINITA DI CONTROLLA DE CONTROL	Lettuce D	'Dallas' (bt:1/28/2005)	
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (include area code)	6. FAX (include area code)	
301 Natividad Road	(831) 443-4901	(831) 443-3976	
Salinas, CA 93906 USA 7. PVPO NUMBER 200200132			
9. Is the applicant (individual or company) a U.S. national or U.S. based of If no, give name of country	ompany?	X YES NO	
10. Is the applicant the original owner?	If no, please answer one of the fo	ollowing:	
a. If original rights to variety were owned by individual(s), is (are) the or	riginal owner(s) a U.S. national(s)?		
	If no, give name of country		
b. If original rights to variety were owned by a company(ies), is(are) the	original owner(s) a U.S. based company	?	
☐ YES ☐ N	If no, give name of country		
11. Additional explanation on ownership (if needed, use reverse for extra sp	pace):		
	•		
	•	•	

### PLEASE NOTE:

Plant variety protection can be afforded only to owners (not licensees) who meet one of the following criteria:

- 1. If the rights to the variety are owned by the original breeder, that person must be a U.S. national of a UPOV member country, or national of a country which affords similar protection to nationals of the U.S. for the same genus and species.
- 2. If the rights to the variety are owned by the company which employed the original breeder(s), the company must be U.S. based, owned by nationals of a UPOV member country, or owned by nationals of a country which affords similar protection to nationals of the U.S. for the same genus and species.
- 3. If the applicant is an owner who is not the original owner, both the original owner and the applicant must meet one of the above criteria.

The original breeder/owner may be the individual or company who directed final breeding. See Section 41(a)(2) of the Plant Variety Protection Act for definition.

According to the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to compete this information collection is estimated to average 10 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call 1-800-245-6340 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.

STD-470-E (07-97) (Destroy previous editions).

Electronic version designed using WordPerfect InForms by USDA-AMS-IMB.